

ON THE THEORETICAL UNDERSTANDING OF INCLUSIVE Λ_b DECAYS

PIETRO COLANGELO

Istituto Nazionale di Fisica Nucleare - Sezione di Bari - via Amendola n.173, I-70126 Bari, Italy

I discuss a QCD sum rule determination of the four-quark operator matrix elements contributing to the Λ_b inclusive decay rates at $\mathcal{O}(m_b^{-3})$. The results suggest that $1/m_b^3$ corrections are not responsible of the observed difference between the lifetimes of Λ_b and B_d .

1 Introduction

The difference between the measured lifetimes of the Λ_b baryon and B_d meson¹:

$$\tau(\Lambda_b^0)/\tau(\bar{B}^0) = 0.78 \pm 0.04 \quad (1)$$

represents an intriguing problem of the present-day heavy quark physics. Indeed, the 20% difference contradicts the expectation that, at the scale of the b quark mass, the spectator model should describe rather accurately the decays of hadrons H_Q containing one heavy quark.

A calculation of the ratio $\tau(\Lambda_b)/\tau(B_d)$ can be attempted using a field theoretical approach developed for the analysis of the inclusive weak decays of the hadrons H_Q ². The method is based on the expansion in powers of m_Q^{-1} , in the framework of the Wilson OPE. The widths are expressed in terms of hadronic matrix elements of high dimensional quark and gluon operators; since such matrix elements can be responsible of the large difference between $\tau(\Lambda_b)$ and $\tau(B_d)$, it is interesting to compute them by nonperturbative methods such as QCD sum rules.

The main aspects of the QCD analysis of the inclusive decays of hadrons H_Q can be summarized considering the transition operator $\hat{T}(Q \rightarrow X_f \rightarrow Q)$ ²:

$$\hat{T} = i \int d^4x T[\mathcal{L}_W(x)\mathcal{L}_W^\dagger(0)] \quad (2)$$

which describes an amplitude with the heavy quark Q having the same momentum in the initial and final state. $\mathcal{L}_W$ is the effective weak Lagrangian governing the decay $Q \rightarrow X_f$. The inclusive width $H_Q \rightarrow X_f$ can be obtained from

$$\Gamma(H_Q \rightarrow X_f) = \frac{2 \text{Im} \langle H_Q | \hat{T} | H_Q \rangle}{2 M_{H_Q}}. \quad (3)$$

The large energy release in the heavy quark decay permits an expansion of $\hat{T}$ in terms of local operators $\mathcal{O}_i$:

$$\hat{T} = \sum_i C_i \mathcal{O}_i \quad (4)$$

with $\mathcal{O}_i$ ordered according to the dimension, and the coefficients C_i containing appropriate inverse powers of the heavy quark mass m_Q . The lowest dimension operator in (4) is $\mathcal{O}_3 = \bar{Q}Q$. The next gauge and Lorentz invariant operator is the $D = 5$ chromomagnetic operator $\mathcal{O}_G = \bar{Q} \frac{g}{2} \sigma_{\mu\nu} G^{\mu\nu} Q$, whose hadronic matrix element

$$\mu_G^2(H_Q) = \frac{\langle H_Q | \bar{Q} \frac{g}{2} \sigma_{\mu\nu} G^{\mu\nu} Q | H_Q \rangle}{2 M_{H_Q}} \quad (5)$$

measures the coupling of the heavy quark spin to the spin of the light degrees of freedom in H_Q , and therefore is responsible of the mass splitting between hadrons belonging to the same s_ℓ multiplet (s_ℓ is the total angular momentum of the light degrees of freedom in H_Q). In the case of b -flavoured hadrons this mass difference has been measured, both for mesons ($M_{B^*} - M_B = 45.7 \pm 0.4 \text{ MeV}$) and Σ_b baryons³ ($M_{\Sigma_b^*} - M_{\Sigma_b} = 56 \pm 16 \text{ MeV}$).

The matrix element of $\bar{Q}Q$ over H_Q can be obtained using the following expansion stemming from the heavy quark equation of motion:

$$\bar{Q}Q = \bar{Q}\gamma^0 Q + \frac{\mathcal{O}_G}{2m_Q^2} - \frac{\mathcal{O}_\pi}{2m_Q^2} + \mathcal{O}(m_Q^{-3}) \quad ; \quad (6)$$

$\mathcal{O}_\pi$ is the kinetic energy operator $\mathcal{O}_\pi = \bar{Q}(i\vec{D})^2 Q$ whose matrix element

$$\mu_\pi^2(H_Q) = \frac{\langle H_Q | \bar{Q}(i\vec{D})^2 Q | H_Q \rangle}{2 M_{H_Q}} \quad (7)$$

measures the average squared momentum of the heavy quark inside H_Q .

At the order $\mathcal{O}(m_Q^{-3})$ four-quark operators appear in (4):

$$\mathcal{O}_6^q = \bar{Q}\Gamma q \bar{q}\Gamma Q \quad (8)$$

with Γ an appropriate combination of Dirac and color matrices.

The resulting expression for the width $\Gamma(H_Q \rightarrow X_f)$ reads:

$$\Gamma(H_Q \rightarrow X_f) = \Gamma_0^f \left[A_0^f + \frac{A_2^f}{m_Q^2} + \frac{A_3^f}{m_Q^3} + \dots \right] \quad (9)$$

A_i^f and Γ_0^f depend on the final state X_f ; A_i^f include perturbative short-distance coefficients and nonperturbative hadronic matrix elements incorporating the long range dynamics. The leading term in (9) corresponds to the partonic prediction $\Gamma^{part}(H_Q \rightarrow X_f) = \Gamma_0^f A_0^f$, with $A_0^f = 1 + c^f \alpha_s/\pi + \mathcal{O}(\alpha_s^2)$ and $\Gamma_0^f \propto m_Q^5$; differences among the widths of the hadrons H_Q emerge at the next to leading order in $1/m_Q$, and are related to the different value of the matrix elements of the operators $\mathcal{O}_i$ of dimension larger than three.

It is important to notice the absence of the first order term m_Q^{-1} in (9)⁴.

The $D = 5$ operators $\mathcal{O}_G$ and $\mathcal{O}_\pi$ are SU(3) singlets; on the contrary, the $D = 6$ operators in (8) give different contributions when averaged over hadrons belonging to the same SU(3) light flavour multiplet, and therefore they are responsible of the different lifetimes of, e.g., B^- and B_s , Λ_b and Ξ_b . The spectator flavour dependence is related to the mechanisms of weak scattering and Pauli interference², both suppressed by the factor m_Q^{-3} with respect to the parton decay rate.

As for differences between mesons and baryons, they could already arise at $\mathcal{O}(m_Q^{-2})$, due both to the chromomagnetic contribution and to the kinetic energy term in (3). In particular, the kinetic energy term is responsible of the difference for systems where the chromomagnetic contribution vanishes, namely in the case of Λ_b and Ξ_b having the light degrees of freedom in S - wave. However, the results of a calculation of μ_π^2 for mesons⁵ and baryons⁶ support the conjecture⁷ that the kinetic energy operator has the same matrix element when computed on such hadronic systems.⁸ The approximate equality of the kinetic energy operator on B_d and Λ_b can also be inferred by using mass relations⁹: $\mu_\pi^2(\Lambda_b) - \mu_\pi^2(B_d) \simeq 0.002 \pm 0.024 \text{ GeV}^2$. Then, differences between

meson and baryon lifetimes should occur at the m_Q^{-3} level, thus involving the four-quark operators in (8). They can be classified as follows¹⁰:

$$\begin{aligned} \mathcal{O}_{V-A}^q &= \bar{Q}_L \gamma_\mu q_L \bar{q}_L \gamma_\mu Q_L \\ \mathcal{O}_{S-P}^q &= \bar{Q}_R q_L \bar{q}_L Q_R \\ T_{V-A}^q &= \bar{Q}_L \gamma_\mu \frac{\lambda^a}{2} q_L \bar{q}_L \gamma_\mu \frac{\lambda^a}{2} Q_L \\ T_{S-P}^q &= \bar{Q}_R \frac{\lambda^a}{2} q_L \bar{q}_L \frac{\lambda^a}{2} Q_R \end{aligned} \quad (10)$$

($q_{R,L} = \frac{1 \pm \gamma_5}{2} q$, λ_a Gell-Mann matrices).

For mesons, the matrix elements of the operators in (10) can be computed by vacuum saturation approximation:

$$\langle B_q | \mathcal{O}_{V-A}^q | B_q \rangle_{VSA} = \frac{f_{B_q}^2 M_{B_q}^2}{4} \quad (11)$$

$$\langle B_q | T_{V-A}^q | B_q \rangle_{VSA} = 0 \quad , \quad (12)$$

etc. Such an approximation cannot be employed for baryons, where a direct calculation is required.

For Λ_b , a simplification can be obtained¹⁰ introducing

$$\tilde{\mathcal{O}}_{V-A}^q = \bar{Q}_L^i \gamma_\mu Q_L^i \bar{q}_L^j \gamma_\mu q_L^j \quad (13)$$

and

$$\tilde{\mathcal{O}}_{S-P}^q = \bar{Q}_L^i q_R^j \bar{q}_L^j Q_R^i \quad (14)$$

(i and j color indices): the Λ_b matrix elements of the operators in (10) can be expressed in terms of $\langle \Lambda_b | \tilde{\mathcal{O}}_{V-A}^q | \Lambda_b \rangle$ and $\langle \Lambda_b | \mathcal{O}_{V-A}^q | \Lambda_b \rangle$, modulo $1/m_Q$ corrections contributing to subleading terms in the expression for the inclusive widths.

Parametrizing the matrix elements

$$\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b} = \frac{\langle \Lambda_b | \tilde{\mathcal{O}}_{V-A}^q | \Lambda_b \rangle}{2M_{\Lambda_b}} = \frac{f_B^2 M_B}{48} r \quad (15)$$

$$\langle \Lambda_b | \mathcal{O}_{V-A}^q | \Lambda_b \rangle = -\tilde{B} \langle \Lambda_b | \tilde{\mathcal{O}}_{V-A}^q | \Lambda_b \rangle \quad , \quad (16)$$

one has that, for $f_B = 200 \text{ MeV}$ and $r = 1$, eq.(15) corresponds to $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b} = 4.4 \times 10^{-3} \text{ GeV}^3$. Quark models predict $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_c} \simeq 0.75 - 2.5 \times 10^{-3} \text{ GeV}^3$, corresponding to $r \simeq 0.2 - 0.6$ ¹¹; larger values can be obtained¹² using the mass splitting $\Sigma_b^* - \Sigma_b$ and $\Sigma_c^* - \Sigma_c$: $r \simeq 0.9 - 1.8$. In the valence quark approximation $\tilde{B} = 1$.

A value of r : $r \simeq 4 - 5$ would explain the difference between $\tau(\Lambda_b)$ and $\tau(B_d)$ ¹⁰. A calculation by QCD sum rules, however, seems to exclude this possibility.

2 $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b}$ from QCD sum rules

An estimate of $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b}$ can be obtained by QCD sum rules, in HQET, analyzing the three-point correlator

$$\begin{aligned} \Pi_{CD}(\omega, \omega') &= (1 + \not{v})_{CD} \Pi(\omega, \omega') \\ &= i^2 \int dx dy \langle 0 | T [J_C(x) \tilde{\mathcal{O}}_{V-A}^q(0) \bar{J}_D(y)] | 0 \rangle \\ &\times e^{i\omega v \cdot x - i\omega' v \cdot y} \end{aligned} \quad (17)$$

of the baryonic currents $J(x)$ and $\bar{J}(y)$ and of the operator $\tilde{\mathcal{O}}_{V-A}^q$ in (13). The variables ω, ω' are related to the residual momentum of the currents: $p^\mu = m_b v^\mu + \omega v^\mu$. Choosing J with a non-vanishing projection on the Λ_b state

$$\langle 0 | J_C | \Lambda_b(v) \rangle = f_{\Lambda_b} (\psi_v)_C \quad (18)$$

(ψ_v is a spinor for a Λ_b of four-velocity v), the parameter f_{Λ_b} representing the coupling of J to Λ_b , the matrix element $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b}$ can be obtained by saturating the correlator (17) with baryonic states, and considering the double pole contribution in the variables ω and ω' :

$$\Pi^h(\omega, \omega') = \langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b} \frac{f_{\Lambda_b}^2}{2} \frac{1}{(\Delta_{\Lambda_b} - \omega)(\Delta_{\Lambda_b} - \omega')} \quad (19)$$

at the value $\omega = \omega' = \Delta_{\Lambda_b}$. The mass parameter Δ_{Λ_b} represents the binding energy of the light Λ_b degrees of freedom in the static color field generated by the b -quark: $M_{\Lambda_b} = m_b + \Delta_{\Lambda_b}$; it must also be derived by QCD sum rules.

A possible interpolating field for Λ_b ¹³ reads:

$$J_C(x) = \epsilon^{ijk} (q^{Ti}(x) \Gamma \tau q^j(x)) (h_v^k)_C(x) \quad (20)$$

where T means transpose, i, j and k are color indices, and C is the Dirac index of the effective heavy quark field $h_v(x)$; τ is a light flavour matrix corresponding to zero isospin. In the $m_b \rightarrow \infty$ limit, the light diquark in the Λ_b is in a relative 0^+ spin-parity state; therefore one can use in (20):

$$\Gamma = C \gamma_5 (1 + b \not{v}) \quad (21)$$

C being the charge conjugation matrix and b a parameter. Arguments can be found in favour of the choice $b = 1$ in (21)⁶.

The coupling f_{Λ_b} can be computed from^{6,14}

$$\begin{aligned} H_{CD}(\omega) &= (1 + \not{v})_{CD} H(\omega) \\ &= i \int dx \langle 0 | T [J_C(x) \bar{J}_D(0)] | 0 \rangle e^{i\omega v \cdot x} \end{aligned} \quad (22)$$

In the Euclidean region the correlation functions (17) and (22) can be computed in QCD, in terms of a perturbative contribution and vacuum condensates:

$$\Pi^{OPE}(\omega, \omega') = \int d\sigma d\sigma' \frac{\rho_\Pi(\sigma, \sigma')}{(\sigma - \omega)(\sigma' - \omega')} \quad (23)$$

$$H^{OPE}(\omega) = \int d\sigma \frac{\rho_H(\sigma)}{(\sigma - \omega)}, \quad (24)$$

with the spectral functions given by :

$$\begin{aligned} \rho_\Pi(\sigma, \sigma') &= \rho_\Pi^{(pert)}(\sigma, \sigma') + \rho_\Pi^{(D=3)}(\sigma, \sigma') \langle \bar{q}q \rangle \\ &+ \rho_\Pi^{(D=4)}(\sigma, \sigma') \langle \frac{\alpha_s}{\pi} G^2 \rangle \\ &+ \rho_\Pi^{(D=5)}(\sigma, \sigma') \langle \bar{q}g\sigma Gq \rangle \\ &+ \rho_\Pi^{(D=6)}(\sigma, \sigma') (\langle \bar{q}q \rangle)^2 + \dots \end{aligned} \quad (25)$$

and a similar expression for $\rho_H(\sigma)$.

The various terms in $\rho_\Pi(\sigma, \sigma')$ and $\rho_H(\sigma)$ can be found in^{6,15}.

A sum rule for $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b}$ can be derived by equating the hadronic and the OPE representations of the correlator (17), and modeling the contribution of the higher resonances and of the continuum in Π^h as the QCD term outside the region $0 \leq \omega \leq \omega_c$, $0 \leq \omega' \leq \omega_c$, ω_c being an effective threshold. After a double Borel transform one gets:

$$\frac{f_{\Lambda_b}^2}{2} (1 + b)^2 e^{-\frac{\Delta_{\Lambda_b}}{E}} \langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b} \quad (26)$$

$$= \int_0^{\omega_c} \int_0^{\omega_c} d\sigma d\sigma' e^{-\frac{\sigma + \sigma'}{2E}} \rho_\Pi(\sigma, \sigma') \quad (27)$$

where E is a Borel parameter. The threshold ω_c can be fixed in the QCD sum rule determination of f_{Λ_b} and Δ_{Λ_b} ⁶: $\omega_c = 1.1 - 1.3 \text{ GeV}$; in correspondence one obtains $f_{\Lambda_b} = (2.9 \pm 0.5) \times 10^{-2} \text{ GeV}^3$ and $\Delta_{\Lambda_b} = 0.9 \pm 0.1 \text{ GeV}$.

The resulting sum rule is depicted in fig.1. A stability window is observed for $E > 0.2 \text{ GeV}$. In the duality region $E \simeq 0.2 - 0.3 \text{ GeV}$ (the same region considered in the QCD sum rule analysis of f_{Λ_b} and $\mu_\pi^2(\Lambda_b)$ ⁶) we find:

$$\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b} \simeq (0.4 - 1.20) \times 10^{-3} \text{ GeV}^3, \quad (28)$$

which corresponds to $r \simeq 0.1 - 0.3$. This result is confirmed by an analysis based on the assumption of local quark-hadron duality, that amounts to calculate the matrix elements of $\langle \tilde{\mathcal{O}}_{V-A}^q \rangle_{\Lambda_b}$ and f_{Λ_b}




Figure 1: Sum rule for the matrix element $\langle \tilde{O}_{V-A}^q \rangle_{\Lambda_b}$

by free quark states produced and annihilated by the baryonic currents in (17) and (22), and then averaging on a duality interval in ω, ω' ¹⁵.

Considering, finally, the parameter $\tilde{B}$ in eq.(16), one gets $\tilde{B} = 1$ since, in this computational scheme, only valence quark processes are taken into account.

3 Conclusions

Within the uncertainties of the method, QCD sum rules predict small values for the matrix elements $\langle \tilde{O}_{V-A}^q \rangle_{\Lambda_b}$ and $\langle O_{V-A}^q \rangle_{\Lambda_b}$, comparable with the outcome of constituent quark models. The conclusion is that the inclusion of $1/m_Q^3$ terms in the expression of the inclusive widths does not solve the puzzle represented by the difference between $\tau(\Lambda_b)$ and $\tau(B_d)$: using the formulae in¹⁰ for the lifetime ratio, the value in (28) together with $\tilde{B} = 1$ gives:

$$\tau(\Lambda_b)/\tau(B_d) \geq 0.94 \quad . \quad (29)$$

It seems unlikely that order m_Q^{-4} terms can solve the problem. We must conclude that, if the measurement of $\tau(\Lambda_b)$ and $\tau(B_d)$ will be confirmed, a reanalysis of the problem will be required. In particular, it has been proposed to consider the possibility of the failure of the assumption, made in the calculation of the lifetimes, of local quark-hadron duality in nonleptonic inclusive decays^{16,17,18}. Meanwhile, it is interesting that new data are now available for other b -flavoured hadrons, e.g. Ξ_b^{-1} , although with errors too large to perform a meaningful comparison with Λ_b . Such new information will be of great importance for the full understanding of the problem of the beauty hadron lifetimes.

Acknowledgments

I would like to thank F.De Fazio, C.A.Dominguez, G.Nardulli and N.Paver for the pleasant collaboration on the topics discussed in this talk.

References

1. J.Richman, these Proceedings.
2. B.Blok and M.Shifman in *Proceedings of the Third Workshop on the Tau-Charm Factory*, eds. J.Kirkby and R.Kirkby, (Ed. Frontieres, 1994) p.269, and references therein.
3. P.Abreu *et al.*, DELPHI Collaboration, preprint DELPHI 95-107 PHYS 542 (1995).
4. J.Chay *et al.*, *Phys. Lett. B* **247**, 399 (1990); I.Bigi *et al.*, *Phys. Lett. B* **293**, 430 (1992); **297**,477 (1993) (E).
5. P.Ball and V.M.Braun, *Phys. Rev. D* **49**, 2472 (1994).
6. P.Colangelo, C.A.Dominguez, G.Nardulli and N.Paver, *Phys. Rev. D* **54**, 4622 (1996).
7. A.V.Manohar and M.B.Wise, *Phys. Rev. D* **49**, 1310 (1994).
8. The value of μ_π^2 is quite controversial: for a discussion see M.Neubert, hep-ph/9610471 and references therein.
9. I.Bigi, hep-ph/9508408.
10. M.Neubert and C.T.Sachrajda, hep-ph/9603202.
11. B.Guberina *et al.*, *Z. Phys. C* **33**, 297 (1986).
12. J.L.Rosner, *Phys. Lett. B* **379**, 267 (1996).
13. E.V.Shuryak, *Nucl. Phys. B* **198**, 83 (1982).
14. A.G.Grozin *et al.*, *Phys. Lett. B* **285**, 254 (1992); **291**, 441 (1992); E.Bagan *et al.*, *Phys. Lett. B* **278**, 367 (1992); **287**, 176 (1992); **301**, 243 (1993); Y.B.Dai *et al.*, *Phys. Lett. B* **371**, 99 (1996); S.Groote *et al.*, *Phys. Rev. D* **54**, 3477 (1996); hep-ph/9609469.
15. P.Colangelo and F.De Fazio, *Phys. Lett. B* **387**, 371 (1996).
16. A.Falk, M.B.Wise and I.Dunietz, *Phys. Rev. D* **51**, 1183 (1995).
17. G.Altarelli *et al.*, *Phys. Lett. B* **382**, 409 (1996).
18. G.Martinelli, these Proceedings.